

## RESEARCH

## Open Access

# On multivalued weakly Picard operators in partial Hausdorff metric spaces

Mohamed Jleli<sup>1</sup>, Hemant Kumar Nashine<sup>2</sup>, Bessem Samet<sup>1\*</sup> and Calogero Vetro<sup>3</sup>

\*Correspondence:

bsamet@ksu.edu.sa

<sup>1</sup>Department of Mathematics,  
College of Science, King Saud  
University, P.O. Box 2455, Riyadh,  
11451, Saudi Arabia

Full list of author information is  
available at the end of the article

## Abstract

We discuss multivalued weakly Picard operators on partial Hausdorff metric spaces. First, we obtain Kikkawa-Suzuki type fixed point theorems for a new type of generalized contractive conditions. Then, we prove data dependence of a fixed points set theorem. Finally, we present sufficient conditions for well-posedness of a fixed point problem. Our results generalize, complement and extend classical theorems in metric and partial metric spaces.

**MSC:** 47H10; 54H25

**Keywords:** data dependence; fixed point; multivalued operator; partial metric space

## 1 Introduction and preliminaries

In 1937, von Neumann [1] initiated the fixed point theory for multivalued mappings in the study of game theory. Indeed, the fixed point theorems for multivalued mappings are quite useful in control theory and have been frequently used in solving many problems of economics. In 1969, Nadler [2] initiated the development of the metric fixed point theory for multivalued mappings. Nadler used the concept of Hausdorff metric to establish the multivalued contraction principle containing the Banach contraction principle as a special case. Also, for the basic problems of fixed point theory for multivalued mappings, we refer to [3].

Let  $(X, d)$  be a metric space and let  $CB(X)$  be the family of all nonempty, closed and bounded subsets of  $X$ . For  $A, B \in CB(X)$ ,  $x \in X$ , let

$$H(A, B) = \max \left\{ \sup_{x \in A} d(x, B), \sup_{y \in B} d(y, A) \right\},$$

where  $d(x, B) = \inf \{d(x, y) : y \in B\}$  and  $H : CB(X) \times CB(X) \rightarrow \mathbb{R}^+$  is the Hausdorff metric induced by  $d$ .

Now on, the letters  $\mathbb{R}$ ,  $\mathbb{R}^+$  and  $\mathbb{N}$  will denote the set of all real numbers, the set of all non-negative real numbers and the set of all positive integers, respectively. Also,  $CL(X)$  is the collection of nonempty closed subsets of  $X$ .

**Definition 1.1** Let  $X$  be a nonempty set. If  $T : X \rightarrow CB(X)$  is a multivalued operator, then an element  $x \in X$  is called

- (i) fixed point of  $T$  if  $x \in Tx$ ;
- (ii) strict fixed point of  $T$  if  $\{x\} = Tx$ .

In the sequel, we denote by  $\text{Fix}(T) := \{x \in X : x \in Tx\}$  the set of all fixed points of  $T$  and by  $S\text{Fix}(T) := \{x \in X : \{x\} = Tx\}$  the set of all strict fixed points of  $T$ .

**Definition 1.2** ([4]) Let  $(X, d)$  be a metric space and  $T : X \rightarrow \text{CL}(X)$  be a multivalued operator.  $T$  is called a multivalued weakly Picard operator (briefly MWP operator) if for all  $x \in X$  and all  $y \in Tx$ , there exists a sequence  $\{x_n\}$  such that:

- (i)  $x_0 = x, x_1 = y$ ;
- (ii)  $x_{n+1} \in Tx_n$  for all  $n \in \mathbb{N} \cup \{0\}$ ;
- (iii) the sequence  $\{x_n\}$  is convergent and its limit is a fixed point of  $T$ .

A sequence  $\{x_n\}$  satisfying (i) and (ii) is also called a sequence of successive approximations (briefly s.s.a.) of  $T$  starting from  $x_0$ .

For interested readers, the theory of MWP operators was presented in [4–7].

In 2008 Suzuki [8] introduced a new type of mappings in order to generalize the well-known Banach contraction principle. This result has led to some important contributions in metric fixed point theory (see, for instance, [9] and the references therein).

As we mentioned above, Nadler proved the following multivalued version of the Banach contraction principle.

**Theorem 1.3** ([2]) Let  $(X, d)$  be a complete metric space and  $T : X \rightarrow \text{CB}(X)$  be a multivalued mapping satisfying  $H(Tx, Ty) \leq kd(x, y)$  for all  $x, y \in X$  and  $k \in (0, 1)$ . Then  $T$  has a fixed point.

In the last decades, a number of fixed point results (see [10–17]) have been obtained in attempts to generalize Theorem 1.3.

One of the most significant fixed point theorems for multivalued mappings appeared in [9], Theorem 2.1. This theorem merges the ideas of Suzuki [8] and Nadler [2] into a consistent framework.

**Theorem 1.4** ([9]) Let  $(X, d)$  be a complete metric space and  $T : X \rightarrow \text{CB}(X)$ . Consider the non-increasing function  $\psi : [0, 1] \rightarrow (0, 1]$  defined by

$$\psi(r) = \begin{cases} 1 & \text{if } 0 \leq r < \frac{1}{2}, \\ 1-r & \text{if } \frac{1}{2} \leq r < 1. \end{cases}$$

Assume that there exists  $r \in [0, 1)$  such that

$$\psi(r)d(x, Tx) \leq d(x, y) \implies H(Tx, Ty) \leq rM_d(x, y)$$

for all  $x, y \in X$ , where

$$M_d(x, y) = \max \left\{ d(x, y), d(x, Tx), d(y, Ty), \frac{d(x, Ty) + d(y, Tx)}{2} \right\}.$$

Then  $T$  has a fixed point.

We remark that the right-hand side of the above implication is known as Ćirić type contractive condition, see [10, 11, 18]. Also, for our further use, we recall the following refinement of Nadler's theorem, see [14].

**Theorem 1.5** Let  $\eta : [0, 1) \rightarrow (\frac{1}{2}, 1]$  be a function defined by  $\eta(r) = \frac{1}{1+r}$ . Let  $(X, d)$  be a complete metric space and  $T : X \rightarrow \text{CB}(X)$  be an  $r$ -KS multivalued operator, that is, there exists  $r \in [0, 1)$  such that

$$\eta(r)d(x, Tx) \leq d(x, y) \implies H(Tx, Ty) \leq rd(x, y)$$

for all  $x, y \in X$ . Then  $T$  is an MWP operator.

The other basic notion for the development of our work is the concept of partial metric space, which was introduced by Matthews [19] as a part of the study of denotational semantics of dataflow networks. Matthews presented a modified version of the Banach contraction principle, more suitable in this context, see also [20, 21]. For more reading on interesting approaches to partial metric spaces and related contexts, we refer to [22–24]. Now, the (complete) partial metric spaces constitute a suitable framework to model several distinguished examples of the theory of computation and also to model metric spaces via domain theory, see [18, 19, 25–37]. In this direction, Aydi *et al.* [38] introduced the concept of partial Hausdorff metric and extended Nadler's fixed point theorem in the setting of partial metric spaces.

Consistent with [19, 38–40], the following definitions and results will be needed in the sequel.

**Definition 1.6** ([19]) Let  $X$  be any nonempty set. A function  $p : X \times X \rightarrow \mathbb{R}^+$  is said to be a partial metric if and only if for all  $x, y, z \in X$  the following conditions are satisfied:

- (P1)  $p(x, x) = p(y, y) = p(x, y)$  if and only if  $x = y$ ;
- (P2)  $p(x, x) \leq p(x, y)$ ;
- (P3)  $p(x, y) = p(y, x)$ ;
- (P4)  $p(x, z) \leq p(x, y) + p(y, z) - p(y, y)$ .

The pair  $(X, p)$  is called a partial metric space. If  $p(x, y) = 0$ , then (P1) and (P2) imply that  $x = y$ , but the converse does not hold in general. A trivial example of a partial metric space is the pair  $(\mathbb{R}^+, p)$ , where  $p : \mathbb{R}^+ \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is given by  $p(x, y) = \max\{x, y\}$ , see also [41].

**Example 1.7** ([19]) Let  $X = \{[a, b] : a, b \in \mathbb{R}, a \leq b\}$ . It is easy to show that the function  $p : X \times X \rightarrow \mathbb{R}^+$  given by  $p([a, b], [c, d]) = \max\{b, d\} - \min\{a, c\}$  defines a partial metric on  $X$ .

For further examples, we refer to [34, 35, 39, 42, 43]. Note that each partial metric  $p$  on  $X$  generates a  $T_0$  topology  $\tau_p$  on  $X$  which has as a base the family of the open balls ( $p$ -balls)  $\{B_p(x, \epsilon) : x \in X, \epsilon > 0\}$ , where for all  $x \in X$  and  $\epsilon > 0$ ,

$$B_p(x, \epsilon) = \{y \in X : p(x, y) < p(x, x) + \epsilon\}.$$

A sequence  $\{x_n\}$  in a partial metric space  $(X, p)$  is called convergent to a point  $x \in X$ , with respect to  $\tau_p$ , if and only if  $p(x, x) = \lim_{n \rightarrow +\infty} p(x, x_n)$ , see [19] for details. If  $p$  is a partial metric on  $X$ , then the function

$$p^S(x, y) = 2p(x, y) - p(x, x) - p(y, y)$$

defines a metric on  $X$ . Further a sequence  $\{x_n\}$  converges in the metric space  $(X, p^S)$  to a point  $x \in X$  if and only if

$$p(x, x) = \lim_{n \rightarrow +\infty} p(x, x_n) = \lim_{n, m \rightarrow +\infty} p(x_n, x_m).$$

**Definition 1.8** ([19]) Let  $(X, p)$  be a partial metric space. Then:

- (i) A sequence  $\{x_n\}$  in  $X$  is called Cauchy if and only if  $\lim_{n, m \rightarrow +\infty} p(x_n, x_m)$  exists and is finite.
- (ii) A partial metric space  $(X, p)$  is said to be complete if every Cauchy sequence  $\{x_n\}$  in  $X$  converges, with respect to  $\tau_p$ , to a point  $x \in X$  such that  $p(x, x) = \lim_{n, m \rightarrow +\infty} p(x_n, x_m)$ .

**Lemma 1.9** ([19, 39]) Let  $(X, p)$  be a partial metric space. Then:

- (i) A sequence  $\{x_n\}$  in  $X$  is Cauchy in  $(X, p)$  if and only if it is Cauchy in  $(X, p^S)$ .
- (ii) A partial metric space  $(X, p)$  is complete if and only if the metric space  $(X, p^S)$  is complete.

Consistent with [38], let  $(X, p)$  be a partial metric space and let  $CB^p(X)$  be the family of all nonempty, closed and bounded subsets of the partial metric space  $(X, p)$ , induced by the partial metric  $p$ . Note that the closedness is taken from  $(X, \tau_p)$  ( $\tau_p$  is the topology induced by  $p$ ) and the boundedness is given as follows:  $A$  is a bounded subset in  $(X, p)$  if there exist  $x_0 \in X$  and  $M \geq 0$  such that for all  $a \in A$ , we have  $a \in B_p(x_0, M)$ , that is,  $p(x_0, a) < p(x_0, x_0) + M$ . For  $A, B \in CB^p(X)$ ,  $x \in X$ ,  $\delta_p : CB^p(X) \times CB^p(X) \rightarrow \mathbb{R}^+$  define

$$\begin{aligned} p(x, A) &= \inf\{p(x, a) : a \in A\}, \\ p(A, B) &= \inf\{p(x, y) : x \in A, y \in B\}, \\ \delta_p(A, B) &= \sup\{p(a, B) : a \in A\}, \\ H_p(A, B) &= \max\{\delta_p(A, B), \delta_p(B, A)\}, \end{aligned}$$

where  $H_p : CB^p(X) \times CB^p(X) \rightarrow \mathbb{R}^+$  is called the partial Hausdorff metric induced by  $p$ . Also, it is easy to show that  $p(x, A) = 0$  implies that  $p^S(x, A) = 0$ , where

$$p^S(x, A) = \inf\{p^S(x, a) : a \in A\}.$$

**Lemma 1.10** ([39]) Let  $(X, p)$  be a partial metric space and  $A$  be any nonempty subset of  $X$ , then  $a \in \overline{A}$  if and only if  $p(a, A) = p(a, a)$ .

**Lemma 1.11** Let  $(X, p)$  be a partial metric space and  $A$  be any nonempty subset of  $X$ . If  $A$  is closed in  $(X, p)$ , then  $A$  is closed in  $(X, p^S)$ .

*Proof* Let  $\{x_n\}$  be a sequence converging to some  $x \in X$  in  $(X, p^S)$ . Then we have

$$p(x, x) = \lim_{n \rightarrow +\infty} p(x, x_n),$$

which implies, by definition, that  $\{x_n\}$  converges to  $x \in X$  also in  $(X, p)$ . Now, since  $A$  is closed in  $(X, p)$ , then  $x \in A$  and so we deduce that  $A$  is closed also in  $(X, p^S)$ .  $\square$

**Proposition 1.12** ([38]) *Let  $(X, p)$  be a partial metric space. For any  $A, B, C \in \text{CB}^p(X)$ , we have:*

- (i)  $\delta_p(A, A) = \sup\{p(a, a) : a \in A\}$ ;
- (ii)  $\delta_p(A, A) \leq \delta_p(A, B)$ ;
- (iii)  $\delta_p(A, B) = 0 \implies A \subseteq B$ ;
- (iv)  $\delta_p(A, B) \leq \delta_p(A, C) + \delta_p(C, B) - \inf_{c \in C} p(c, c)$ .

**Proposition 1.13** ([38]) *Let  $(X, p)$  be a partial metric space. For any  $A, B, C \in \text{CB}^p(X)$ , we have:*

- (h1)  $H_p(A, A) \leq H_p(A, B)$ ;
- (h2)  $H_p(A, B) = H_p(B, A)$ ;
- (h3)  $H_p(A, B) \leq H_p(A, C) + H_p(C, B) - \inf_{c \in C} p(c, c)$ ;
- (h4)  $H_p(A, B) = 0 \implies A = B$ .

Notice that each Hausdorff metric is a partial Hausdorff metric but the converse is not true, see Example 2.6 in [38].

**Lemma 1.14** ([38]) *Let  $(X, p)$  be a partial metric space,  $A, B \in \text{CB}^p(X)$  and  $q > 1$ , then for any  $a \in A$ , there exists  $b(a) \in B$  such that  $p(a, b(a)) \leq qH_p(A, B)$ .*

**Theorem 1.15** ([38]) *Let  $(X, p)$  be a partial metric space. If  $T : X \rightarrow \text{CB}^p(X)$  is a multivalued mapping such that for all  $x, y \in X$ , we have  $H_p(Tx, Ty) \leq kp(x, y)$ , where  $k \in (0, 1)$ , then  $T$  has a fixed point.*

In view of the above considerations and following the ideas in [44], the aim of this paper is to discuss multivalued weakly Picard operators on partial Hausdorff metric spaces, see also [45] for other interesting results. First, we obtain Kikkawa-Suzuki type fixed point theorems for a new type of generalized contractive conditions. Then, we prove data dependence of a fixed points set theorem. Finally, we present sufficient conditions for well-posedness of a fixed point problem. The presented results extend and unify some recently obtained comparable results for multivalued mappings (see [9] and the references therein).

## 2 Fixed point theorems in partial Hausdorff metric spaces

In this section we present several theorems which characterize MWP operators, defined in the previous section, in terms of different contractive conditions. Results of this section are generalizations of Theorem 1.4, Theorem 1.15 (and so Nadler's Theorem 1.3), Ćirić's theorem in [10] and others.

### 2.1 Result - I

To provide the first theorem we introduce the notion of  $(s, r)$ -contractive multivalued operator in partial Hausdorff metric spaces as follows.

**Definition 2.1** Let  $p : X \times X \rightarrow \mathbb{R}^+$  be a partial metric and  $T : X \rightarrow \text{CB}^p(X)$  be a multivalued mapping.  $T$  is called an  $(s, r)$ -contractive multivalued operator if there exist  $r \in [0, 1)$  and  $s \geq r$  such that

$$p(y, Tx) \leq sp(y, x) \implies H_p(Tx, Ty) \leq rM_p(x, y) \quad (1)$$

for all  $x, y \in X$ , where

$$M_p(x, y) = \max \left\{ p(x, y), p(x, Tx), p(y, Ty), \frac{p(x, Ty) + p(y, Tx)}{2} \right\}.$$

Now we state and prove our theorem.

**Theorem 2.2** *Let  $(X, p)$  be a complete partial metric space and  $T : X \rightarrow \text{CB}^p(X)$  be an  $(s, r)$ -contractive multivalued operator with  $s \geq 1$ . Then  $T$  is an MWP operator.*

*Proof* Let  $r_1$  be a real number such that  $0 \leq r < r_1 < 1$ . Let  $u_1 \in X$ . As  $Tu_1$  is nonempty, we can choose  $u_2 \in Tu_1$ . Clearly, if  $u_2 = u_1$  the proof is finished and so we assume  $u_2 \neq u_1$ . Then we get

$$p(u_2, Tu_1) = p(u_2, u_2) \leq p(u_2, u_1) \leq sp(u_2, u_1). \quad (2)$$

Next, we choose  $u_3 \in Tu_2$  such that

$$p(u_3, u_2) \leq \frac{r_1}{r} H_p(Tu_2, Tu_1).$$

Now, from (2) and by using condition (1) we write

$$\begin{aligned} p(u_3, u_2) &\leq \frac{r_1}{r} H_p(Tu_2, Tu_1) \leq r_1 M_p(u_1, u_2) \\ &= r_1 \max \left\{ p(u_1, u_2), p(u_1, Tu_1), p(u_2, Tu_2), \frac{p(u_1, Tu_2) + p(u_2, Tu_1)}{2} \right\} \\ &\leq r_1 \max \left\{ p(u_1, u_2), p(u_1, u_2), p(u_2, u_3), \frac{p(u_1, u_3) + p(u_2, u_2)}{2} \right\} \\ &\leq r_1 \max \left\{ p(u_1, u_2), p(u_2, u_3), \frac{p(u_1, u_2) + p(u_2, u_3)}{2} \right\} \\ &\leq r_1 \max \{ p(u_1, u_2), p(u_2, u_3) \}. \end{aligned}$$

Thus, we obtain

$$p(u_2, u_3) \leq r_1 \max \{ p(u_1, u_2), p(u_2, u_3) \}.$$

If  $\max \{ p(u_1, u_2), p(u_2, u_3) \} = p(u_2, u_3)$ , then  $p(u_2, u_3) \leq r_1 p(u_2, u_3)$  implies that  $p(u_2, u_3) = 0$ , and we obtain  $p^S(u_2, u_3) \leq 2p(u_2, u_3) = 0$ , which further implies that  $p^S(u_2, u_3) = 0$ . Hence  $u_2 = u_3 \in Tu_2$  and the proof is finished. On the contrary, if  $\max \{ p(u_1, u_2), p(u_2, u_3) \} = p(u_1, u_2)$ , then we have

$$p(u_2, u_3) \leq r_1 p(u_1, u_2).$$

Continuing this process, we can construct a sequence  $\{u_n\}$  in  $X$  such that  $u_{n+1} \in Tu_n$ ,  $u_{n+1} \neq u_n$  and

$$p(u_n, u_{n+1}) \leq r_1^{n-1} p(u_1, u_2)$$

for every  $n > 1$ . This shows that  $\lim_{n \rightarrow +\infty} p(u_n, u_{n+1}) = 0$ .

Now, since

$$p(u_n, u_n) + p(u_{n+1}, u_{n+1}) \leq 2p(u_n, u_{n+1}),$$

then we get

$$\lim_{n \rightarrow +\infty} p(u_n, u_n) = 0 \quad \text{and} \quad \lim_{n \rightarrow +\infty} p(u_{n+1}, u_{n+1}) = 0. \quad (3)$$

Let  $\epsilon > 0$  and pick  $N \in \mathbb{N}$  large enough so that for  $n \geq N$  we have

$$2r_1^{n-1} \frac{1}{1-r_1} p(u_1, u_2) < \epsilon.$$

Then, for every positive integer  $k > n \geq N$ , there is some  $m \in \mathbb{N}$  such that  $k = n + m$ , and we have

$$\begin{aligned} p^S(u_n, u_{n+m}) &\leq 2p(u_n, u_{n+m}) \\ &\leq 2[p(u_n, u_{n+1}) + \cdots + p(u_{n+m-1}, u_{n+m})] \\ &\leq 2[r_1^{n-1} p(u_1, u_2) + \cdots + r_1^{n+m-2} p(u_1, u_2)] \\ &\leq 2r_1^{n-1} \frac{1}{1-r_1} p(u_1, u_2) < \epsilon. \end{aligned}$$

It is immediate to deduce that  $\{u_n\}$  is a Cauchy sequence in  $(X, p^S)$ , but by Lemma 1.9  $\{u_n\}$  is Cauchy also in  $(X, p)$ . Moreover, since  $(X, p)$  is complete, again by Lemma 1.9 we deduce the completeness of  $(X, p^S)$ . It follows that there exists  $z \in X$  such that  $\lim_{n \rightarrow +\infty} u_n = z$  in  $(X, p^S)$ . Therefore  $\lim_{n \rightarrow +\infty} p^S(u_n, z) = 0$  implies

$$p(z, z) = \lim_{n \rightarrow +\infty} p(u_n, z) = \lim_{m, n \rightarrow +\infty} p(u_n, u_m).$$

Now, since  $\{u_n\}$  is a Cauchy sequence in  $(X, p^S)$ , then we have

$$\lim_{m, n \rightarrow +\infty} p^S(u_n, u_m) = 0$$

and so

$$\lim_{m, n \rightarrow +\infty} 2p(u_n, u_m) - \lim_{m \rightarrow +\infty} p(u_m, u_m) - \lim_{n \rightarrow +\infty} p(u_n, u_n) = 0.$$

It follows from (3) that

$$\lim_{m \rightarrow +\infty} p(u_m, u_m) = \lim_{n \rightarrow +\infty} p(u_n, u_n) = 0,$$

which further implies that

$$\lim_{m, n \rightarrow +\infty} p(u_n, u_m) = 0$$

and hence

$$p(z, z) = \lim_{n \rightarrow +\infty} p(u_n, z) = \lim_{n \rightarrow +\infty} p(u_n, u_n) = 0.$$

Next, we will show that there exists a subsequence  $\{u_{n(k)}\}$  of  $\{u_n\}$  such that

$$p(z, Tu_{n(k)}) \leq sp(z, u_{n(k)})$$

for all  $k \in \mathbb{N}$ . Reasoning by contradiction, we assume that there exists a positive integer  $N$  such that  $p(z, Tu_n) > sp(z, u_n)$  for all  $n \geq N$ . This implies  $p(z, u_{n+1}) > sp(z, u_n)$  for all  $n \geq N$ . By induction, for all  $n \geq N$  and  $m' \geq 1$ , we get that

$$p(z, u_{n+m'}) > s^{m'} p(z, u_n). \quad (4)$$

Since

$$\begin{aligned} p(u_{n+m'}, u_n) &\leq p(u_n, u_{n+1}) + p(u_{n+1}, u_{n+2}) + \cdots + p(u_{n+m'-1}, u_{n+m'}) - \sum_{i=n+1}^{m'-1} p(u_i, u_i) \\ &\leq p(u_n, u_{n+1}) + p(u_{n+1}, u_{n+2}) + \cdots + p(u_{n+m'-1}, u_{n+m'}) \\ &\leq p(u_n, u_{n+1})(1 + r_1 + r_1^2 + r_1^3 + \cdots + r_1^{m'-1}) \\ &\leq p(u_n, u_{n+1}) \left[ \frac{1 - r_1^{m'}}{1 - r_1} \right] \end{aligned}$$

for all  $n \geq N$  and  $m' \geq 1$ , then we get

$$\begin{aligned} p(z, u_n) &\leq p(z, u_{n+m'}) + p(u_{n+m'}, u_n) - p(z, z) \\ &\leq p(z, u_{n+m'}) + p(u_n, u_{n+1}) \left[ \frac{1 - r_1^{m'}}{1 - r_1} \right]. \end{aligned}$$

Passing to the limit as  $m' \rightarrow +\infty$ , we have

$$p(z, u_n) \leq \frac{1}{1 - r_1} p(u_n, u_{n+1}) \quad \text{for all } n \geq N.$$

Then we obtain

$$p(z, u_{n+m'}) \leq \frac{1}{1 - r_1} p(u_{n+m'}, u_{n+m'+1}) \leq \frac{r_1^{m'}}{1 - r_1} p(u_n, u_{n+1}) \quad (5)$$

for all  $n \geq N$  and  $m' \geq 1$ . By (4) and (5) we get

$$p(z, u_n) < \frac{\left(\frac{r_1}{s}\right)^{m'}}{1 - r_1} p(u_n, u_{n+1})$$

for all  $n \geq N$  and  $m' \geq 1$ . Next, passing to the limit as  $m' \rightarrow +\infty$ , we obtain that  $p(z, u_n) = 0$  for all  $n \geq N$ . This contradicts (4) and therefore there exists a subsequence  $\{u_{n(k)}\}$  of  $\{u_n\}$  such that  $p(z, Tu_{n(k)}) \leq sp(z, u_{n(k)})$  for all  $k \in \mathbb{N}$ .



Also, by hypothesis we have

$$\begin{aligned} H_p(Tu_{n(k)}, Tz) &\leq r \max \left\{ p(u_{n(k)}, z), p(u_{n(k)}, Tu_{n(k)}), p(z, Tz), \frac{p(u_{n(k)}, Tz) + p(z, Tu_{n(k)})}{2} \right\} \\ &\leq r \max \left\{ p(u_{n(k)}, z), p(u_{n(k)}, u_{n(k)+1}), p(z, Tz), \frac{p(u_{n(k)}, z) + p(z, Tz) + p(z, u_{n(k)+1})}{2} \right\}. \end{aligned}$$

Therefore, we write

$$\begin{aligned} p(z, Tz) &\leq p(z, u_{n(k)+1}) + H_p(Tu_{n(k)}, Tz) - p(u_{n(k)+1}, u_{n(k)+1}) \\ &\leq p(z, u_{n(k)+1}) \\ &\quad + r \max \left\{ p(u_{n(k)}, z), p(u_{n(k)}, u_{n(k)+1}), p(z, Tz), \frac{p(u_{n(k)}, z) + p(z, Tz) + p(z, u_{n(k)+1})}{2} \right\}. \end{aligned}$$

On passing to the limit as  $k \rightarrow +\infty$ , we get

$$p(z, Tz) \leq r \max \left\{ p(z, Tz), \frac{p(z, Tz)}{2} \right\} = rp(z, Tz).$$

Since  $r < 1$ , it follows that

$$p(z, Tz) = 0.$$

Therefore  $p(z, Tz) = 0 = p(z, z)$  and hence by Lemma 1.10 we deduce that  $z \in Tz$ , that is,  $z$  is a fixed point of  $T$ .  $\square$

The following example illustrates the use of Theorem 2.2.

**Example 2.3** Let  $X = \{0, 1, 2\}$  and  $p : X \times X \rightarrow \mathbb{R}^+$  be defined by  $p(0, 0) = 0$ ,  $p(1, 1) = p(2, 2) = \frac{1}{4}$ ,  $p(1, 0) = \frac{1}{3}$ ,  $p(2, 0) = \frac{3}{5}$ ,  $p(2, 1) = \frac{2}{5}$  and  $p(y, x) = p(x, y)$  for all  $x, y \in X$ . Then  $(X, p)$  is a complete partial metric space. Also define  $T : X \rightarrow CB^p(X)$  by

$$Tx = \begin{cases} \{0\} & \text{if } x \neq 2, \\ \{0, 1\} & \text{if } x = 2. \end{cases}$$

Therefore, we get

$$\begin{aligned} H_p(T0, T0) &= H_p(T0, T1) = H_p(T1, T1) = p(0, T0) = p(0, T1) = p(0, 0), \\ H_p(T0, T2) &= H_p(T1, T2) = p(1, T0) = p(1, T1) = p(1, 0), \\ p(2, T2) &= \min\{p(2, 0), p(2, 1)\} = p(2, 1), \\ H_p(T2, T2) &= p(2, 2). \end{aligned}$$

It follows easily that the inequality

$$p(y, Tx) \leq sp(x, y)$$

holds for all  $x, y \in X$  with  $s \geq \frac{8}{5}$ . Also, for all  $x, y \in X$  with  $r \in [\frac{5}{6}, 1)$ , we get

$$H_p(Tx, Ty) \leq rM_p(x, y).$$

Thus all the conditions of Theorem 2.2 are satisfied and 0 is a fixed point of  $T$ .

From Theorem 2.2 we deduce some corollaries.

**Corollary 2.4** *Let  $(X, p)$  be a complete partial metric space and  $T : X \rightarrow \text{CB}^p(X)$  be a multivalued mapping. Assume that there exist  $r \in [0, 1)$  and  $s \geq 1$  such that*

$$p(y, Tx) \leq sp(x, y) \implies H_p(Tx, Ty) \leq r \max\{p(x, y), p(x, Tx), p(y, Ty)\}$$

*for all  $x, y \in X$ . Then  $T$  is an MWP operator.*

**Corollary 2.5** *Let  $(X, p)$  be a complete partial metric space and  $T : X \rightarrow \text{CB}^p(X)$  be a multivalued mapping. Assume that there exist  $r \in [0, 1)$  and  $s \geq 1$  such that*

$$p(y, Tx) \leq sp(x, y) \implies H_p(Tx, Ty) \leq \frac{r}{3} [p(x, y) + p(x, Tx) + p(y, Ty)]$$

*for all  $x, y \in X$ . Then  $T$  is an MWP operator.*

In the case of single-valued mappings, Theorem 2.2 reduces to the following significant corollary.

**Corollary 2.6** *Let  $(X, p)$  be a complete partial metric space and  $T : X \rightarrow X$  be a single-valued mapping. Assume that there exist  $r \in [0, 1)$  and  $s \geq 1$  such that*

$$p(y, Tx) \leq sp(x, y) \implies p(Tx, Ty) \leq rM_p(x, y), \quad (6)$$

*where*

$$M_p(x, y) = \max \left\{ p(x, y), p(x, Tx), p(y, Ty), \frac{p(x, Ty) + p(y, Tx)}{2} \right\}.$$

*Then  $T$  has a unique fixed point.*

*Proof* The existence part of the proof follows easily by Theorem 2.2. Thus, we need to prove uniqueness of the fixed point. Suppose to the contrary that, for  $s \geq 1$ , there exist  $x, y \in \text{Fix}(T)$  with  $x \neq y$ . It follows immediately that

$$p(y, Tx) = p(y, x) \leq sp(y, x).$$

Thus, by hypothesis on  $T$ , we would have

$$\begin{aligned} p(Tx, Ty) &\leq r \max \left\{ p(x, y), p(x, Tx), p(y, Ty), \frac{p(x, Ty) + p(y, Tx)}{2} \right\} \\ &= r \max \left\{ p(x, y), p(x, x), p(y, y), \frac{p(x, y) + p(y, x)}{2} \right\} \\ &= rp(x, y). \end{aligned}$$

We deduce that  $p(x, y) = 0$ , which further implies that  $p^S(x, y) \leq 2p(x, y) = 0$  and hence  $x = y$ , a contradiction. This completes the proof.  $\square$

The following two examples, adapted from [46], show the validity of Corollary 2.6.

**Example 2.7** Let  $X = [0, 1]$  be endowed with the partial metric  $p(x, y) = \max\{x, y\}$  for all  $x, y \in X$ . Then  $(X, p)$  is a complete partial metric space. Also define  $T : X \rightarrow X$  by

$$Tx = \frac{x^2}{1+x}.$$

Take arbitrary elements  $x, y \in X$  with  $y \leq x$ . Then we have

$$p(y, Tx) = \max \left\{ y, \frac{x^2}{1+x} \right\} \leq sx = sp(x, y)$$

for all  $x, y \in X$  and  $s \geq 1$ . On the other hand, we get

$$p(Tx, Ty) = \max \left\{ \frac{x^2}{1+x}, \frac{y^2}{1+y} \right\} = \frac{x^2}{1+x}$$

and

$$\begin{aligned} M_p(x, y) &= \max \left\{ p(x, y), p(x, Tx), p(y, Ty), \frac{p(x, Ty) + p(y, Tx)}{2} \right\} \\ &= \max \left\{ p(x, y), p\left(x, \frac{x^2}{1+x}\right), p\left(y, \frac{y^2}{1+y}\right), \frac{1}{2} \left( p\left(x, \frac{y^2}{1+y}\right) + p\left(y, \frac{x^2}{1+x}\right) \right) \right\} \\ &= \max \left\{ x, x, y, \frac{1}{2} \left( x + \max \left\{ y, \frac{x^2}{1+x} \right\} \right) \right\} = x. \end{aligned}$$

Thus the inequality  $p(Tx, Ty) \leq rM_p(x, y)$  holds for all  $x, y \in X$  with  $y \leq x$  and for any  $r \in [\frac{1}{2}, 1)$ . Note that we obtain the same conclusion if we assume that  $x \leq y$ . Hence, all the conditions of Corollary 2.6 are satisfied and 0 is a fixed point of  $T$ .

The following example underlines the crucial role of the right-hand side of (6) in establishing existence of the fixed point.

**Example 2.8** Let  $X = \{1, 2, 3, 4\}$  and  $p : X \times X \rightarrow \mathbb{R}^+$  be defined by  $p(x, x) = \frac{1}{2}$  for each  $x \in X$ ,  $p(1, 2) = p(3, 4) = 2$ ,  $p(1, 3) = p(2, 4) = 1$ ,  $p(1, 4) = p(2, 3) = \frac{3}{2}$  and  $p(y, x) = p(x, y)$  for

all  $x, y \in X$ . Then  $(X, p)$  is a complete partial metric space. Also define  $T : X \rightarrow X$  by

$$T = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 1 & 1 & 2 \end{pmatrix}.$$

Trivially  $T$  has no fixed points, but we try to apply Corollary 2.6. It is easy to check that the inequality  $p(y, Tx) \leq sp(y, x)$  certainly holds for all  $x, y \in X$  with  $s \geq 2$ . Now we note that, for  $x = 3$  and  $y = 1$ , we get

$$p(T3, T1) = p(1, 2) = 2$$

and

$$\begin{aligned} M_p(3, 1) &= \max \left\{ p(3, 1), p(3, T3), p(1, T1), \frac{p(3, T1) + p(1, T3)}{2} \right\} \\ &= \max \left\{ 1, 1, 2, \frac{1}{2} \left( \frac{3}{2} + \frac{1}{2} \right) \right\} = 2. \end{aligned}$$

It follows that

$$p(T3, T1) = 2 \not\leq 2r = rM_p(3, 1),$$

whatever  $r \in [0, 1)$  is chosen. We conclude that Corollary 2.6 is not applicable in this case.

## 2.2 Result - II

Another interesting characterization of MWP operators is provided by the following theorem.

**Theorem 2.9** *Let  $(X, p)$  be a complete partial metric space and  $T : X \rightarrow CB^p(X)$  be a multivalued operator. Assume that there exist  $r, s \in [0, 1)$ , with  $r < s$ , such that*

$$\frac{1}{1+r}p(x, Tx) \leq p(x, y) \leq \frac{1}{1-s}p(x, Tx) \implies H_p(Tx, Ty) \leq rM_p(x, y) \quad (7)$$

for all  $x, y \in X$ , where

$$M_p(x, y) = \max \left\{ p(x, y), p(x, Tx), p(y, Ty), \frac{p(x, Ty) + p(y, Tx)}{2} \right\}.$$

Then  $T$  is an MWP operator.

*Proof* Let  $r_1$  be a real number such that  $0 \leq r < r_1 < s$ . Also, let  $u_1 \in X$  and  $u_2 \in Tu_1$  be such that

$$p(u_1, u_2) \leq \frac{1-r_1}{1-s}p(u_1, Tu_1).$$

Then we have

$$\frac{1}{1+r}p(u_1, Tu_1) \leq p(u_1, Tu_1) \leq p(u_1, u_2) \leq \frac{1}{1-s}p(u_1, Tu_1)$$

and so, by using condition (7), we obtain

$$\begin{aligned} p(u_2, Tu_2) &\leq H_p(Tu_1, Tu_2) \\ &\leq r \max \left\{ p(u_1, u_2), p(u_1, Tu_1), p(u_2, Tu_2), \frac{p(u_1, Tu_2) + p(u_2, Tu_1)}{2} \right\} \\ &\leq r \max \left\{ p(u_1, u_2), p(u_2, Tu_2), \frac{p(u_1, Tu_2) + p(u_2, u_2)}{2} \right\} \\ &\leq r \max \left\{ p(u_1, u_2), p(u_2, Tu_2), \frac{p(u_1, u_2) + p(u_2, Tu_2)}{2} \right\}. \end{aligned}$$

Thus

$$p(u_2, Tu_2) \leq r \max \{p(u_1, u_2), p(u_2, Tu_2)\}.$$

Now, if  $\max\{p(u_1, u_2), p(u_2, Tu_2)\} = p(u_2, Tu_2)$ , then  $p(u_2, Tu_2) \leq rp(u_2, Tu_2)$  implies that  $p(u_2, Tu_2) = 0$  and so we obtain

$$p^S(u_2, Tu_2) \leq 2p(u_2, Tu_2) = 0,$$

which further implies that  $p^S(u_2, Tu_2) = 0$ . In view of Lemma 1.11,  $u_2 \in Tu_2$ , and the proof is finished. On the contrary, if  $\max\{p(u_1, u_2), p(u_2, u_3)\} = p(u_1, u_2)$ , then we have

$$p(u_2, Tu_2) \leq rp(u_1, u_2).$$

It follows that there exists  $u_3 \in Tu_2$  such that

$$p(u_2, u_3) \leq r_1 p(u_1, u_2) \quad \text{and} \quad p(u_2, u_3) \leq \frac{1-r_1}{1-s} p(u_2, Tu_2).$$

This implies

$$\frac{1}{1+r} p(u_2, Tu_2) \leq p(u_2, u_3) \leq \frac{1}{1-s} p(u_2, Tu_2).$$

Now, by using condition (7), we get  $p(u_3, Tu_3) \leq rp(u_2, u_3)$ . Continuing this process, we can construct a sequence  $\{u_n\}$  in  $X$  with the following properties:

$$\begin{cases} u_{n+1} \in Tu_n, \\ p(u_{n+1}, Tu_{n+1}) \leq rp(u_n, u_{n+1}), \\ p(u_{n+1}, u_{n+2}) \leq r_1 p(u_n, u_{n+1}), \\ p(u_{n+1}, u_{n+2}) \leq \frac{1-r_1}{1-s} p(u_{n+1}, Tu_{n+1}) \end{cases}$$

for every  $n \in \mathbb{N}$ . Next, from  $p(u_{n+1}, u_{n+2}) \leq r_1 p(u_n, u_{n+1})$  we deduce that

$$\lim_{n \rightarrow +\infty} p(u_n, u_{n+1}) = 0.$$

Since

$$p(u_n, u_n) + p(u_{n+1}, u_{n+1}) \leq 2p(u_n, u_{n+1}),$$

then we get

$$\lim_{n \rightarrow +\infty} p(u_n, u_n) = 0 \quad \text{and} \quad \lim_{n \rightarrow +\infty} p(u_{n+1}, u_{n+1}) = 0.$$

Proceeding as in the proof of Theorem 2.2, one can show that the sequence  $\{u_n\}$  is a Cauchy sequence in  $(X, p)$  converging to some  $z \in X$  with  $p(z, z) = 0$ .

Now, since

$$p(u_n, u_{n+1}) \leq \frac{1-r_1}{1-s} p(u_n, Tu_n),$$

then we have

$$p(z, u_n) \leq \frac{1}{1-s} p(u_n, Tu_n)$$

for all  $n \geq 1$ . Then we assume that there exists a positive integer  $N$  such that

$$p(z, u_n) < \frac{1}{1+r} p(u_n, Tu_n)$$

holds for every  $n \geq N$ . Consequently, we have

$$\begin{aligned} p(u_n, u_{n+1}) &\leq p(z, u_n) + p(z, u_{n+1}) - p(z, z) \\ &< \frac{1}{1+r} [p(u_n, Tu_n) + p(u_{n+1}, Tu_{n+1})] \\ &< \frac{1}{1+r} [p(u_n, Tu_n) + rp(u_n, u_{n+1})] \\ &\leq \frac{1}{1+r} [p(u_n, u_{n+1}) + rp(u_n, u_{n+1})] = p(u_n, u_{n+1}), \end{aligned}$$

which is a contradiction. Hence, there exists a subsequence  $\{u_{n(k)}\}$  of  $\{u_n\}$  such that

$$p(z, u_{n(k)}) \geq \frac{1}{1+r} p(u_{n(k)}, Tu_{n(k)})$$

holds for every  $k \geq N$ . Since  $p(z, u_n) \leq \frac{1}{1-s} p(u_n, Tu_n)$  for all  $n \geq 1$ , by condition (7), we have  $H_p(Tz, Tu_{n(k)}) \leq rM_p(z, u_{n(k)})$ . This implies

$$\begin{aligned} p(z, Tz) &\leq p(u_{n(k)+1}, z) + H_p(Tu_{n(k)}, Tz) \\ &\leq p(u_{n(k)+1}, z) \\ &\quad + r \max \left\{ p(u_{n(k)}, z), p(u_{n(k)}, Tu_{n(k)}), p(z, Tz), \frac{p(u_{n(k)}, Tz) + p(z, Tu_{n(k)})}{2} \right\} \\ &\leq r \max \left\{ p(u_{n(k)}, z), p(u_{n(k)}, u_{n(k)+1}), p(z, Tz), \frac{p(u_{n(k)}, Tz) + p(z, u_{n(k)+1})}{2} \right\}. \end{aligned}$$

On passing to the limit as  $k \rightarrow +\infty$ , we get

$$p(z, Tz) \leq r \max \left\{ p(z, Tz), \frac{p(z, Tz)}{2} \right\}.$$

Since  $r < 1$ , it follows that

$$p(z, Tz) = 0.$$

Therefore  $p(z, Tz) = 0 = p(z, z)$  and hence by Lemma 1.10 we have  $z \in Tz$ , that is,  $z$  is a fixed point of  $T$ .  $\square$

In the case of single-valued mappings, Theorem 2.9 reduces to the following corollary.

**Corollary 2.10** *Let  $(X, p)$  be a complete partial metric space and  $T : X \rightarrow X$  be a single-valued mapping. Assume that there exists  $r \in [0, 1)$  such that*

$$\frac{1}{1+r}p(x, Tx) \leq p(x, y) \leq \frac{1}{1-r}p(x, Tx) \implies p(Tx, Ty) \leq rM_p(x, y)$$

for all  $x, y \in X$ , where

$$M_p(x, y) = \max \left\{ p(x, y), p(x, Tx), p(y, Ty), \frac{p(x, Ty) + p(y, Tx)}{2} \right\}.$$

Then  $T$  has a fixed point.

*Proof* It is easy to prove that for every  $u_1 \in X$  the sequence  $\{u_n\}$  defined by  $u_{n+1} = Tu_n$  satisfies the relationship  $p(u_{n+1}, u_{n+2}) \leq rp(u_n, u_{n+1})$ . Consequently, the sequence  $\{u_n\}$  is Cauchy, and there is some point  $z \in X$  such that  $\lim_{n \rightarrow +\infty} u_n = z$ . Proceeding as in the proof of Theorem 2.9, we can show that

$$p(z, z) = \lim_{n \rightarrow +\infty} p(u_n, z) = \lim_{n \rightarrow +\infty} p(u_n, u_n) = 0.$$

Also in view of Theorem 2.9, for all  $n \geq N$ , we can assume  $p(z, u_n) \leq \frac{1}{1+r}p(u_n, u_{n+1})$  for leading to contradiction. Consequently, there exists a subsequence  $\{u_{n(k)}\}$  of  $\{u_n\}$  such that

$$p(z, u_{n(k)}) \geq \frac{1}{1+r}p(u_{n(k)}, u_{n(k)+1})$$

holds for every  $k \geq N$ . Therefore, we obtain that

$$\begin{aligned} & p(u_{n(k)+1}, Tz) \\ & \leq r \max \left\{ p(u_{n(k)}, z), p(u_{n(k)}, Tu_{n(k)}), p(z, Tz), \frac{p(u_{n(k)}, Tz) + p(z, Tu_{n(k)})}{2} \right\} \\ & \leq r \max \left\{ p(u_{n(k)}, z), p(u_{n(k)}, u_{n(k)+1}), p(z, Tz), \frac{p(u_{n(k)}, z) + p(z, Tz) + p(z, u_{n(k)+1})}{2} \right\}. \end{aligned}$$

On passing to the limit as  $k \rightarrow +\infty$ , we get

$$p(z, Tz) \leq r \max \left\{ p(z, Tz), \frac{p(z, Tz)}{2} \right\}.$$

Since  $r < 1$ , it follows that

$$p(z, Tz) = 0$$

and so  $z = Tz$ , that is,  $z$  is a fixed point of  $T$ .  $\square$

The following example shows that Theorem 2.9 is proper extension of the respective result in standard metric spaces.

**Example 2.11** Let  $X = \{0, 1, 2\}$  and  $p : X \times X \rightarrow \mathbb{R}^+$  be defined by  $p(0, 0) = p(1, 1) = 0$ ,  $p(2, 2) = \frac{1}{4}$ ,  $p(1, 0) = \frac{1}{3}$ ,  $p(2, 0) = \frac{2}{5}$ ,  $p(2, 1) = \frac{11}{15}$  and  $p(y, x) = p(x, y)$  for all  $x, y \in X$ . Then  $(X, p)$  is a complete partial metric space. Also define  $T : X \rightarrow CB^p(X)$  by

$$Tx = \begin{cases} \{0\} & \text{if } x \neq 2, \\ \{0, 1\} & \text{if } x = 2. \end{cases}$$

Therefore, we get

$$\min\{p(x, Tx) : x \in X \setminus \{0\}\} = \min\{p(x, y) : x, y \in X \text{ and } x \neq y\} = \frac{1}{3}$$

and

$$\max\{p(x, Tx) : x \in X\} = \frac{2}{5}.$$

It follows easily that the inequalities

$$\frac{1}{1+r}p(x, Tx) \leq p(x, y) \leq \frac{1}{1-s}p(x, Tx)$$

hold for all  $x, y \in X$  with  $x \neq y$  and for some  $1 > s > r \geq \frac{1}{5}$ . Also the above inequalities hold for  $x = y = 2$  with  $1 > s > r \geq \frac{3}{5}$ . On the other hand, the above inequalities are not applicable for  $x = y \in \{0, 1\}$ . Clearly we have

$$H_p(T0, T0) = H_p(T0, T1) = H_p(T1, T1) = p(0, T0) = p(0, T1) = p(0, 0),$$

$$H_p(T0, T2) = H_p(T1, T2) = p(1, T0) = p(1, T1) = p(1, 0),$$

$$p(2, T2) = \min\{p(2, 0), p(2, 1)\} = p(2, 0),$$

$$H_p(T2, T2) = p(2, 2).$$

Finally, by routine calculations and taking  $1 > s > r \geq \frac{5}{6}$ , one can show that the inequality

$$H_p(Tx, Ty) \leq rM_p(x, y)$$

holds true as for all  $x, y \in X$  with  $x \neq y$ , as for  $x = y = 2$ . Thus all the conditions of Theorem 2.9 are satisfied and 0 is a fixed point of  $T$ .



Next, we consider the metric  $p^S$  induced by the partial metric  $p$ . Indeed, we have  $p^S(x, x) = 0$  for all  $x \in X$ ,  $p^S(1, 0) = \frac{2}{3}$ ,  $p^S(2, 1) = \frac{73}{60}$ ,  $p^S(2, 0) = \frac{11}{20}$  and  $p^S(x, y) = p^S(y, x)$  for all  $x, y \in X$ .

We show that Theorem 2.9 is not applicable in this case. Indeed, for  $x = 2$  and  $y = 0$ , the inequalities

$$\frac{1}{1+r}p^S(2, T2) = \frac{1}{1+r}p^S(2, \{0, 1\}) = \frac{1}{1+r}p^S(2, 0) \leq p^S(2, 0) \leq \frac{1}{1-s}p^S(2, T2)$$

hold true for all  $r \in [0, 1)$  with  $r < s$ . Therefore, we need to have

$$H_{p^S}(T0, T2) \leq rM_{p^S}(0, 2).$$

Unfortunately, this is not the case because

$$H_{p^S}(T0, T2) = H_{p^S}(\{0\}, \{0, 1\}) = \frac{2}{3}$$

and

$$\begin{aligned} M_{p^S}(0, 2) &= \max \left\{ p^S(0, 2), p^S(0, T0), p^S(2, T2), \frac{p^S(0, T2) + p^S(2, T0)}{2} \right\} \\ &= \max \left\{ \frac{11}{20}, 0, \frac{11}{20}, \frac{0 + \frac{11}{20}}{2} \right\} = \frac{11}{20} < \frac{2}{3}. \end{aligned}$$

Consequently, for any  $r \in [0, 1)$  we have

$$H_{p^S}(T0, T2) \not\leq rM_{p^S}(0, 2).$$

### 3 Data dependence theorem in partial Hausdorff metric spaces

The aim of this section is to discuss data dependence of a fixed points set for MWP operators on partial metric spaces. Also, this section is motivated by Popescu [44], see also [4]. Precisely, we will prove a result for  $(1, r)$ -contractive multivalued operators in partial Hausdorff metric spaces.

First, we need the following auxiliary lemma.

**Lemma 3.1** *Let  $(X, p)$  be a partial metric space and  $T : X \rightarrow \text{CB}^p(X)$  be a  $(1, r)$ -contractive multivalued operator. If  $z \in Tz$ , then  $p(z, z) = 0$ .*

*Proof* Since  $p(z, Tz) = p(z, z)$ , then we have

$$H_p(Tz, Tz) \leq r \max \left\{ p(z, z), p(z, Tz), p(z, Tz), \frac{p(z, Tz) + p(z, Tz)}{2} \right\} = rp(z, z).$$

Thus, by definition, we write

$$p(z, z) \leq p(z, Tz) \leq H_p(Tz, Tz) \leq rp(z, z) < p(z, z)$$

and hence we deduce that  $p(z, z) = 0$ . □

We recall the following concept.

**Definition 3.2** ([47]) Let  $(X, p)$  be a partial metric space and let  $\phi : X \rightarrow \mathbb{R}^+$  be a function on  $X$ . Then the function  $\phi$  is called  $p$ -lower semi-continuous on  $X$  whenever

$$\lim_{n \rightarrow +\infty} p(x_n, x) = p(x, x) \implies \phi(x) \leq \lim_{n \rightarrow +\infty} \inf \phi(x_n) = \sup_{n \geq 1} \inf_{m \geq n} \phi(x_m).$$

Now we state and prove our theorem.

**Theorem 3.3** Let  $(X, p)$  be a partial metric space and  $T_1, T_2 : X \rightarrow \text{CB}^p(X)$  be two multi-valued operators. We suppose that:

- (i)  $T_i$  is a  $(1, r_i)$ -contractive multivalued operator for  $i \in \{1, 2\}$ ;
- (ii) there exists  $\lambda > 0$  such that  $H_p(T_1x, T_2x) \leq \lambda$  for all  $x \in X$ ;
- (iii) the function  $\phi : X \rightarrow \mathbb{R}^+$  defined by  $\phi(x) = p(x, x)$  is  $p$ -lower semi-continuous.

Then:

- (a)  $\text{Fix}(T_i) \in \text{CL}^p(X)$ ,  $i \in \{1, 2\}$ ;
- (b)  $T_1$  and  $T_2$  are MWP operators and

$$H_p(\text{Fix}(T_1), \text{Fix}(T_2)) \leq \frac{\lambda}{1 - \max\{r_1, r_2\}}.$$

*Proof* (a) From Theorem 2.2 we have that  $\text{Fix}(T_i)$  is a nonempty set,  $i \in \{1, 2\}$ . Let us prove that the fixed point set of a  $(1, r_i)$ -contractive multivalued operator  $T_i$  is closed. Let  $x_n \in \text{Fix}(T_i)$ , with  $n \geq 1$ , be such that  $\lim_{n \rightarrow +\infty} x_n = z$  in  $(X, p)$ . In view of (iii) and Lemma 3.1, we have

$$\lim_{n \rightarrow +\infty} p(x_n, z) = p(z, z) \implies p(z, z) \leq \lim_{n \rightarrow +\infty} \inf p(x_n, x_n) = 0.$$

It follows that

$$p(z, z) = \lim_{n \rightarrow +\infty} p(x_n, z) = \lim_{n \rightarrow +\infty} p(x_n, x_n) = 0.$$

Also, since  $x_n \in T_i x_n$ , we have  $p(z, T_i x_n) \leq p(z, x_n)$  and then

$$\begin{aligned} p(z, T_i z) &\leq p(z, x_n) + p(x_n, T_i z) - p(x_n, x_n) \\ &\leq p(z, x_n) + H_p(T_i x_n, T_i z) \\ &\leq p(z, x_n) + r_i p(x_n, z). \end{aligned}$$

Passing to the limit as  $n \rightarrow +\infty$ , we obtain that  $p(z, T_i z) = 0$ . Therefore

$$p(z, T_i z) = 0 = p(z, z)$$

and hence by Lemma 1.10 and  $T_i z \in \text{CL}^p(X)$  we get  $z \in T_i z$ , that is,  $z \in \text{Fix}(T_i)$ .

(b) From the proof of Theorem 2.2 we immediately get that a  $(1, r_i)$ -contractive multivalued operator is an MWP operator. For the second conclusion, let  $q$  be a real number such that  $q > 1$ , and  $x_0 \in \text{Fix}(T_1)$  be arbitrary. Then, by Lemma 1.14, there exists  $x_1 \in T_2 x_0$  such that

$$p(x_0, x_1) \leq q H_p(T_1 x_0, T_2 x_0).$$

Next, for  $x_1 \in T_2x_0$ , there exists  $x_2 \in T_2x_1$  such that

$$p(x_1, x_2) \leq qH_p(T_2x_0, T_2x_1).$$

Since  $x_1 \in T_2x_0$ ,  $p(x_1, T_2x_0) = p(x_1, x_1) \leq p(x_0, x_1)$ , then we have

$$p(x_1, x_2) \leq qH_p(T_2x_0, T_2x_1) \leq qr_2p(x_0, x_1).$$

Iterating this process allows us to construct a sequence of successive approximations for  $T_2$  starting from  $x_0$ , satisfying the following assertions:

$$x_{n+1} \in T_2x_n \quad \text{and} \quad p(x_n, x_{n+1}) \leq (qr_2)^n p(x_0, x_1) \quad \text{for all } n \in \mathbb{N}.$$

Hence, for all  $n \geq N$  and  $m \geq 1$ , we write

$$\begin{aligned} p(x_{n+m}, x_n) &\leq p(x_n, x_{n+1}) + p(x_{n+1}, x_{n+2}) + \cdots + p(x_{n+m-1}, x_{n+m}) - \sum_{i=n+1}^{m-1} p(x_i, x_i) \\ &\leq p(x_n, x_{n+1}) + p(x_{n+1}, x_{n+2}) + \cdots + p(x_{n+m-1}, x_{n+m}) \\ &\leq \frac{(qr_2)^n}{1 - qr_2} p(x_0, x_1). \end{aligned}$$

Consequently, we get

$$p^S(x_n, x_{n+m}) \leq 2p(x_n, x_{n+m}) \leq \frac{2(qr_2)^n}{1 - qr_2} p(x_0, x_1). \quad (8)$$

Now, choosing  $1 < q < \min\{\frac{1}{r_1}, \frac{1}{r_2}\}$  and passing to the limit as  $n \rightarrow +\infty$ , we deduce easily that the sequence  $\{x_n\}$  is Cauchy in  $(X, p^S)$  and, by Lemma 1.9,  $\{x_n\}$  is Cauchy in  $(X, p)$ . Then there exists  $u \in X$  such that

$$\lim_{n \rightarrow +\infty} x_n = u \quad \text{in } (X, p^S).$$

Therefore, (8) and  $\lim_{n \rightarrow +\infty} p^S(x_n, u) = 0$  imply

$$p(u, u) = \lim_{n \rightarrow +\infty} p(x_n, u) = \lim_{n, m \rightarrow +\infty} p(x_n, x_m) = 0.$$

We will prove that  $u$  is a fixed point for  $T_2$ . To this aim, suppose that there exists a positive integer  $N$  such that

$$p(u, T_2x_n) > p(u, x_n) \quad \text{for all } n \geq N.$$

This implies that  $p(u, x_{n+1}) > p(u, x_n)$  for all  $n \geq N$ , which leads to contradiction since  $x_n \rightarrow u$  as  $n \rightarrow +\infty$ . Hence, there exists a subsequence  $\{x_{n(k)}\}$  such that

$$p(u, T_2x_{n(k)}) \leq p(u, x_{n(k)}) \quad \text{for all } k \in \mathbb{N}.$$

Next, from

$$p(u, T_2 u) \leq p(x_{n(k)+1}, u) + H_p(T_2 x_{n(k)}, T_2 u),$$

on passing to the limit as  $k \rightarrow +\infty$ , we get

$$\begin{aligned} p(u, T_2 u) &\leq \lim_{k \rightarrow +\infty} p(x_{n(k)+1}, u) + \lim_{k \rightarrow +\infty} H_p(T_2 x_{n(k)}, T_2 u) \\ &\leq \lim_{k \rightarrow +\infty} r_2 p(x_{n(k)}, u) = 0 \end{aligned}$$

and so  $u \in \text{Fix}(T_2)$ .

By (8), passing to the limit as  $m \rightarrow +\infty$ , we get

$$\begin{aligned} p(x_n, u) &\leq \lim_{m \rightarrow +\infty} [p(x_n, x_{n+m}) + p(x_{n+m}, u) - p(u, u)] \\ &\leq \lim_{m \rightarrow +\infty} \left[ \frac{2(qr_2)^n}{1 - qr_2} p(x_0, x_1) + p(x_{n+m}, u) \right] \\ &\leq \frac{(qr_2)^n}{1 - qr_2} p(x_0, x_1) \quad \text{for each } n \in \mathbb{N}. \end{aligned}$$

Then

$$p(x_0, u) \leq \frac{1}{1 - qr_2} p(x_0, x_1) \leq \frac{q\lambda}{1 - qr_2}.$$

Analogously, one can show that, for each  $u_0 \in \text{Fix}(T_2)$ , there exists  $x \in \text{Fix}(T_1)$  such that

$$p(u_0, x) \leq \frac{1}{1 - qr_1} p(u_0, u_1) \leq \frac{q\lambda}{1 - qr_1}$$

and hence

$$H_p(\text{Fix}(T_1), \text{Fix}(T_2)) \leq \frac{q\lambda}{1 - \max\{qr_1, qr_2\}}.$$

Finally, passing to the limit as  $q \rightarrow 1^+$ , we obtain the assertion. This concludes the proof.  $\square$

#### 4 Well-posedness of fixed point problems in partial Hausdorff metric spaces

According to [48, 49], we get the notions of well-posedness of a fixed point problem in the setting of partial metric spaces.

**Definition 4.1** (see [48, 49]) Let  $(X, p)$  be a partial metric space and  $T : X \rightarrow \text{CB}^p(X)$  be a multivalued operator. Then the fixed point problem is well posed for  $T$  with respect to  $p$  if:

- (a<sub>1</sub>)  $\text{Fix}(T) = \{z\}$ ;
- (b<sub>1</sub>) if  $x_n \in X$ ,  $n \in \mathbb{N}$  and  $\lim_{n \rightarrow +\infty} p(x_n, Tx_n) = 0$ , then  $\lim_{n \rightarrow +\infty} p(x_n, z) = 0$ .

**Definition 4.2** (see [48, 49]) Let  $(X, p)$  be a partial metric space and  $T : X \rightarrow \text{CB}^p(X)$  be a multivalued operator. Then the fixed point problem is well posed for  $T$  with respect to  $H_p$  if:

- (a<sub>2</sub>)  $S\text{Fix}(T) = \{z\}$ ;  
 (b<sub>2</sub>) if  $x_n \in X$ ,  $n \in \mathbb{N}$  and  $\lim_{n \rightarrow +\infty} H_p(x_n, Tx_n) = 0$ , then  $\lim_{n \rightarrow +\infty} p(x_n, z) = 0$ .

Clearly, (b<sub>2</sub>) of Definition 4.2 implies (b<sub>1</sub>) of Definition 4.1. Moreover, from (a<sub>1</sub>) and (a<sub>2</sub>), that is,  $\text{Fix}(T) = S\text{Fix}(T) = \{z\}$ , we deduce that if the fixed point problem is well posed for  $T$  with respect to  $p$ , then it is well posed for  $T$  with respect to  $H_p$ .

Motivated by the above facts, we will prove the following theorem for  $(s, r)$ -contractive multivalued operators, with  $s > 1$ , in partial metric spaces.

**Theorem 4.3** *Let  $(X, p)$  be a partial metric space and  $T : X \rightarrow \text{CB}^p(X)$  be a multivalued operator. We suppose that:*

- (1)  *$T$  is an  $(s, r)$ -contractive multivalued operator with  $s \geq 1$ ;*
- (2)  *$S\text{Fix}(T) \neq \emptyset$ .*

*Then:*

- (a)  $\text{Fix}(T) = S\text{Fix}(T) = \{z\}$ ;
- (b) *the fixed point problem is well posed for  $T$  with respect to  $H_p$  if  $s > 1$ .*

*Proof* (a) Suppose  $z \in S\text{Fix}(T)$ . Clearly,  $z \in \text{Fix}(T)$ . We show that  $\text{Fix}(T) = \{z\}$ . To this aim, let  $u \in \text{Fix}(T)$  with  $u \neq z$ . Since  $p(u, Tz) = p(u, z) \leq sp(u, z)$ , we get  $H_p(Tz, Tu) \leq rp(z, u)$  and therefore

$$p(z, u) = p(Tz, u) \leq H_p(Tz, Tu) \leq rp(z, u),$$

which leads to contradiction. Thus,  $\text{Fix}(T) = \{z\}$  and so the assertion (a) holds true.

(b) Now, let  $x_n \in X$ , with  $n \in \mathbb{N}$ , be such that  $\lim_{n \rightarrow +\infty} p(x_n, Tx_n) = 0$ . We have to show that  $\lim_{n \rightarrow +\infty} p(x_n, z) = 0$ . Suppose this is not the case; suppose that  $p(x_n, z)$  does not converge to zero. Consequently, there exist  $\epsilon > 0$  and a subsequence  $\{x_{n(k)}\}$  such that

$$p(x_{n(k)}, z) \geq \epsilon \quad \text{for all } k \in \mathbb{N}.$$

Now, assume that there exists a subsequence  $\{x_{n(k(j))}\}$  of  $\{x_{n(k)}\}$  with

$$p(z, Tx_{n(k(j))}) \leq sp(z, x_{n(k(j))}).$$

Then we get

$$H_p(Tz, Tx_{n(k(j))}) \leq rp(z, x_{n(k(j))}),$$

and so we write

$$\begin{aligned} p(z, x_{n(k(j))}) &= p(x_{n(k(j))}, Tz) \\ &\leq p(x_{n(k(j))}, Tx_{n(k(j))}) + H_p(Tz, Tx_{n(k(j))}) \\ &\leq p(x_{n(k(j))}, Tx_{n(k(j))}) + rp(z, x_{n(k(j))}). \end{aligned}$$

The above inequality leads to the following:

$$\epsilon \leq p(z, x_{n(k(j))}) \leq \frac{1}{1-r} p(x_{n(k(j))}, Tx_{n(k(j))}).$$

Passing to the limit as  $j \rightarrow +\infty$ , since  $\lim_{n \rightarrow +\infty} p(x_n, Tx_n) = 0$ , we get the contradiction  $\epsilon = 0$ . Consequently, we deduce that there exists  $k_1 \in \mathbb{N}$  such that

$$p(z, Tx_{n(k)}) > sp(z, x_{n(k)}) \quad \text{for all } k \geq k_1.$$

Again, since  $\lim_{n \rightarrow +\infty} p(x_n, Tx_n) = 0$ , there exists  $k_2 \geq k_1$  such that

$$p(x_{n(k)}, Tx_{n(k)}) < (s-1)\epsilon \quad \text{for all } k \geq k_2.$$

Finally, for all  $k \geq k_2$ , we write

$$\begin{aligned} (s-1)\epsilon &\leq (s-1)p(z, x_{n(k)}) \\ &= sp(z, x_{n(k)}) - p(z, x_{n(k)}) \\ &< p(z, Tx_{n(k)}) - p(z, x_{n(k)}) \\ &\leq p(x_{n(k)}, Tx_{n(k)}) \quad (\text{by (P4) of Definition 1.6}) \\ &< (s-1)\epsilon, \end{aligned}$$

which leads to contradiction. Consequently, we conclude that  $\lim_{n \rightarrow +\infty} p(x_n, z) = 0$ .  $\square$

## 5 Application to integral equations

The literature is rich with papers focusing on the study of integral operators of various types: Fredholm, Urysohn, Volterra and others. It is well known that integral operators provide an important subject of numerous mathematical investigations and are often applicable in many scientific disciplines as physics, biology and economics. The papers we refer to essentially present a fixed point approach based on the Banach contraction principle and its constructive proof. These results give sufficient conditions for establishing the existence (and uniqueness) of solution of certain integral operators, see [50–52].

Here, following this line of research, we prove an existence theorem for the solution of integral equations by using Corollary 2.6. Precisely, we consider the following integral equation:

$$u(t) = \int_0^T K(t, s, u(s)) ds + g(t), \quad (9)$$

$t \in I = [0, T]$ , where  $T > 0$ . Also we denote

$$C(I) := \{u : I \rightarrow \mathbb{R} \mid u \text{ is continuous on } I\},$$

and define  $d : C(I) \times C(I) \rightarrow \mathbb{R}$  by

$$d(u, v) = \max_{t \in I} |u(t) - v(t)|$$

for all  $u, v \in C(I)$ , so that  $(C(I), d)$  is a complete metric space.

Finally, we define the operator  $T : C(I) \rightarrow C(I)$  by

$$Tx(t) = \int_0^T K(t, s, x(s)) ds + g(t)$$

for all  $x \in C(I)$  and  $t \in I$ .

Then we consider the following hypotheses:

- (i)  $K : I \times I \times \mathbb{R} \rightarrow \mathbb{R}$  and  $g : I \rightarrow \mathbb{R}$  are continuous;
- (ii) there exist  $\alpha \in [0, 1)$ ,  $\beta \geq 1$  and a continuous function  $G : I \times I \rightarrow \mathbb{R}^+$  such that

$$\max_{t \in I} \int_0^T G(t, s) ds = \alpha,$$

and the inequality

$$\max_{t \in I} \left| \int_0^T K(t, s, u(s)) ds + g(t) - v(t) \right| \leq \beta \max_{t \in I} |u(t) - v(t)|$$

implies

$$\begin{aligned} & |K(t, s, u(t)) - K(t, s, v(t))| \\ & \leq G(t, s) \max \left\{ |u(t) - v(t)|, \right. \\ & \quad \left| \int_0^T K(t, s, u(s)) ds + g(t) - u(t) \right|, \left| \int_0^T K(t, s, v(s)) ds + g(t) - v(t) \right|, \\ & \quad \left. \frac{1}{2} \left( \left| \int_0^T K(t, s, v(s)) ds + g(t) - u(t) \right| + \left| \int_0^T K(t, s, u(s)) ds + g(t) - v(t) \right| \right) \right\} \end{aligned}$$

for all  $u, v \in C(I)$  and  $t, s \in I$ .

We will prove the following result.

**Theorem 5.1** *Suppose that hypotheses (i) and (ii) hold. Then the integral equation (9) has a unique solution  $x^* \in C(I)$ .*

*Proof* First we note that the space  $(C(I), d)$  is trivially a complete partial metric space with zero self-distance.

Next, for all  $u, v \in C(I)$ , by (ii), we deduce that

$$\begin{aligned} & |Tu(t) - Tv(t)| \\ & \leq \int_0^T |K(t, s, u(s)) - K(t, s, v(s))| ds \\ & \leq \int_0^T G(t, s) ds \max \left\{ |u(t) - v(t)|, |u(t) - T(u(t))|, \right. \\ & \quad \left. |v(t) - T(v(t))|, \frac{|u(t) - T(v(t))| + |v(t) - T(u(t))|}{2} \right\}, \end{aligned}$$

which, on routine calculations, leads to

$$d(Tu, Tv) \leq \alpha M_d(u, v).$$

Therefore, without loss of generality, we can write that

$$d(v, Tu) \leq \beta d(u, v) \implies d(Tu, Tv) \leq \alpha M_d(u, v)$$

for all  $u, v \in C(I)$ . Thus Corollary 2.6 is applicable in this case, and hence the operator  $T$  has a unique fixed point  $x^* \in C(I)$ . Clearly,  $x^* \in C(I)$  is the unique solution of (9).  $\square$

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript.

#### Author details

<sup>1</sup>Department of Mathematics, College of Science, King Saud University, P.O. Box 2455, Riyadh, 11451, Saudi Arabia.

<sup>2</sup>Department of Mathematics, Disha Institute of Management and Technology, Satya Vihar, Vidhansabha-Chandrakhuri Marg, Mandir Hasaud, Raipur, Chhattisgarh 492101, India. <sup>3</sup>Dipartimento di Matematica e Informatica, Università degli Studi di Palermo, Via Archirafi 34, Palermo, 90123, Italy.

#### Authors' information

CV is a member of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

#### Acknowledgements

The authors would like to extend their sincere appreciation to the Deanship of Scientific Research at King Saud University for its funding of this research through the International Research Group Project No. IRG14-04.

Received: 7 November 2014 Accepted: 27 January 2015 Published online: 11 April 2015

#### References

1. von Neuman, J: Über ein ökonomisches Gleichungssystem und eine Verallgemeinerung des Brouwerschen Fixpunktsatzes. *Ergeb. Math. Kolloqu.* **8**, 73-83 (1937)
2. Nadler, SB Jr: Multi-valued contraction mappings. *Pac. J. Math.* **30**, 475-488 (1969)
3. Petruşel, A, Rus, IA, Şerban, MA: Basic problems of the metric fixed point theory and the relevance of a metric fixed point theorem for a multivalued operator. *J. Nonlinear Convex Anal.* **15**, 493-513 (2014)
4. Rus, IA, Petruşel, A, Sintămărian, A: Data dependence of the fixed points set of multivalued weakly Picard operators. *Stud. Univ. Babeş-Bolyai, Math.* **46**, 111-121 (2001)
5. Moş, G, Petruşel, A: Fixed point theory for a new type of contractive multivalued operators. *Nonlinear Anal.* **70**, 3371-3377 (2009)
6. Petruşel, A: Multivalued weakly Picard operators and applications. *Sci. Math. Jpn.* **59**, 169-202 (2004)
7. Rus, IA, Petruşel, A, Sintămărian, A: Data dependence of the fixed points set of some multivalued weakly Picard operators. *Nonlinear Anal.* **52**, 1947-1959 (2003)
8. Suzuki, T: A generalized Banach contraction principle that characterizes metric completeness. *Proc. Am. Math. Soc.* **136**, 1861-1869 (2008)
9. Dorić, D, Lazović, R: Some Suzuki type fixed point theorems for generalized multivalued mappings and applications. *Fixed Point Theory Appl.* **2011**, 40 (2011)
10. Ćirić, L: Fixed points for generalized multi-valued contractions. *Mat. Vesn.* **9**, 265-272 (1972)
11. Ćirić, L: Multi-valued nonlinear contraction mappings. *Nonlinear Anal.* **71**, 2716-2723 (2009)
12. Covitz, H, Nadler, SB Jr: Multi-valued contraction mappings in generalized metric spaces. *Isr. J. Math.* **8**, 5-11 (1970)
13. Daffer, PZ, Kaneko, H: Fixed points of generalized contractive multi-valued mappings. *J. Math. Anal. Appl.* **192**, 655-666 (1995)
14. Kikkawa, M, Suzuki, T: Some similarity between contractions and Kannan mappings. *Fixed Point Theory Appl.* **2008**, Article ID 649749 (2008)
15. Kikkawa, M, Suzuki, T: Three fixed point theorems for generalized contractions with constants in complete metric spaces. *Nonlinear Anal.* **69**, 2942-2949 (2008)
16. Mizoguchi, N, Takahashi, W: Fixed point theorems for multi-valued mappings on complete metric spaces. *J. Math. Anal. Appl.* **141**, 177-188 (1989)
17. Semenov, PV: Fixed points of multi-valued contractions. *Funct. Anal. Appl.* **36**, 159-161 (2002)
18. Ćirić, L, Samet, B, Aydi, H, Vetro, C: Common fixed points of generalized contractions on partial metric spaces and an application. *Appl. Math. Comput.* **218**, 2398-2406 (2011)
19. Matthews, SG: Partial metric topology. In: *Proc. 8th Summer Conference on General Topology and Applications*. Ann. New York Acad. Sci., vol. 728, pp. 183-197 (1994)
20. Di Bari, C, Vetro, P: Fixed points for weak  $\varphi$ -contractions on partial metric spaces. *Int. J. Eng. Contemp. Math. Sci.* **1**, 5-13 (2011)
21. Paesano, D, Vetro, P: Suzuki's type characterizations of completeness for partial metric spaces and fixed points for partially ordered metric spaces. *Topol. Appl.* **159**, 911-920 (2012)
22. Amini-Harandi, A: Metric-like spaces, partial metric spaces and fixed points. *Fixed Point Theory Appl.* **2012**, 204 (2012)
23. Haghi, RH, Rezapour, S, Shahzad, N: Be careful on partial metric fixed point results. *Topol. Appl.* **160**, 450-454 (2013)
24. Samet, B: Existence and uniqueness of solutions to a system of functional equations and applications to partial metric spaces. *Fixed Point Theory* **14**, 473-482 (2013)
25. Abbas, M, Ali, B, Vetro, C: A Suzuki type fixed point theorem for a generalized multivalued mapping on partial Hausdorff metric spaces. *Topol. Appl.* **160**, 553-563 (2013)
26. Di Bari, C, Kadelburg, Z, Nashine, HK, Radenović, S: Common fixed points of  $g$ -quasicontractions and related mappings in 0-complete partial metric spaces. *Fixed Point Theory Appl.* **2012**, 113 (2012)



27. Di Bari, C, Milojević, M, Radenović, S, Vetro, P: Common fixed points for self-mappings on partial metric spaces. *Fixed Point Theory Appl.* **2012**, 140 (2012)
28. Dukić, D, Kadelburg, Z, Radenović, S: Fixed point of Geraghty-type mappings in various generalized metric spaces. *Abstr. Appl. Anal.* **2011**, Article ID 561245 (2011)
29. Heckmann, R: Approximation of metric spaces by partial metric spaces. *Appl. Categ. Struct.* **7**, 71-83 (1999)
30. Nashine, HK, Karapinar, E: Fixed point results for orbitally continuous map in orbitally complete partial metric spaces. *Bull. Malays. Math. Soc.* **36**, 1185-1193 (2013)
31. Nashine, HK, Kadelburg, Z, Radenović, S: Common fixed point theorems for weakly isotone increasing mappings in ordered partial metric spaces. *Math. Comput. Model.* **57**, 2355-2365 (2013)
32. Nashine, HK, Kadelburg, Z, Radenović, S: Fixed point theorems via various cyclic contractive conditions in partial metric spaces. *Publ. Inst. Math. (Belgr.)* **93**, 69-93 (2013)
33. Nashine, HK, Kadelburg, Z, Radenović, S, Kim, JK: Fixed point theorems under Hardy-Rogers contractive conditions on 0-complete ordered partial metric spaces. *Fixed Point Theory Appl.* **2012**, 180 (2012)
34. Romaguera, S: A Kirk type characterization of completeness for partial metric spaces. *Fixed Point Theory Appl.* **2010**, Article ID 493298 (2010)
35. Schellekens, MP: The correspondence between partial metrics and semivaluations. *Theor. Comput. Sci.* **315**, 135-149 (2004)
36. Vetro, F, Radenović, S: Nonlinear  $\psi$ -quasi-contractions of Ćirić-type in partial metric spaces. *Appl. Math. Comput.* **219**, 1594-1600 (2012)
37. Nashine, HK: Fixed point results for mappings satisfying  $(\phi, \varphi)$ -weakly contractive condition in ordered partial metric spaces. *Math. Slovaca* **63**, 871-882 (2013)
38. Aydi, H, Abbas, M, Vetro, C: Partial Hausdorff metric and Nadler's fixed point theorem on partial metric spaces. *Topol. Appl.* **159**, 3234-3242 (2012)
39. Altun, I, Simsek, H: Some fixed point theorems on dualistic partial metric spaces. *J. Adv. Math. Stud.* **1**, 1-8 (2008)
40. Altun, I, Sola, F, Simsek, H: Generalized contractions on partial metric spaces. *Topol. Appl.* **157**, 2778-2785 (2010)
41. Abbas, M, Nazir, T: Fixed point of generalized weakly contractive mappings in ordered partial metric spaces. *Fixed Point Theory Appl.* **2012**, 1 (2012)
42. Bukatin, MA, Shorina, SY: Partial metrics and co-continuous valuations. In: Nivat, M (ed.) *Foundations of Software Science and Computation Structure. Lect. Notes Comput. Sci.*, vol. 1378, pp. 125-139. Springer, Berlin (1998)
43. Romaguera, S, Valero, O: A quantitative computational model for complete partial metric spaces via formal balls. *Math. Struct. Comput. Sci.* **19**, 541-563 (2009)
44. Popescu, O: A new type of contractive multivalued operators. *Bull. Sci. Math.* **137**, 30-44 (2013)
45. Abbas, M, Ali, B, Petruşel, G: Fixed points of set-valued contractions in partial metric spaces endowed with a graph. *Carpath. J. Math.* **30**, 129-137 (2014)
46. Ahmad, AGB, Fadail, ZM, Nashine, HK, Kadelburg, Z, Radenović, S: Some new common fixed point results through generalized altering distances on partial metric spaces. *Fixed Point Theory Appl.* **2012**, 120 (2012)
47. Karapinar, E: Generalizations of Caristi Kirk's theorem on partial metric spaces. *Fixed Point Theory Appl.* **2011**, 4 (2011)
48. Petruşel, A, Rus, IA: Well-posedness of the fixed point problem for multivalued operators. In: Cârjă, O, Vrabie, II (eds.) *Applied Analysis and Differential Equations*, pp. 295-306. World Scientific, Singapore (2007)
49. Petruşel, A, Rus, IA, Yao, J-C: Well-posedness in the generalized sense of the fixed point problems for multivalued operators. *Taiwan. J. Math.* **11**, 903-914 (2007)
50. Berinde, V: *Iterative Approximation of Fixed Points. Lecture Notes in Math.*, vol. 1912. Springer, Berlin (2007)
51. Granas, A, Dugundji, J: *Fixed Point Theory. Monogr. Math.*, vol. XVI. Springer, Berlin (2003)
52. Jachymski, J, Reich, S: *Fixed Point Theory and Its Applications. Banach Center Publ.* vol. 77. Polish Academy of Sciences, Institute of Mathematics, Warsaw (2007)

**Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:**

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)